



## Improving postcryopreservation survival capacity: an embryo-focused approach

M.J. Sudano<sup>1,4</sup>, D.M. Paschoal<sup>1</sup>, R.R.D. Maziero<sup>1</sup>, T.S. Rascado<sup>1</sup>, M.D. Guastali<sup>1</sup>, L.F. Crocomo<sup>1</sup>, L.C.O. Magalhães<sup>1</sup>, B.A. Monteiro<sup>1</sup>, A. Martins Jr<sup>2</sup>, R. Machado<sup>3</sup>, F.D.C. Landim-Alvarenga<sup>1,4</sup>

<sup>1</sup>São Paulo State University, School of Veterinary Medicine and Animal Science, Department of Animal Reproduction and Veterinary Radiology, Botucatu, SP, Brazil.

<sup>2</sup>São Paulo State University, School of Veterinary Medicine, Department of Clinical Surgery and Animal Reproduction, Araçatuba, SP, Brazil.

<sup>3</sup>Embrapa Southeast Cattle, São Carlos, SP, Brazil.

### Abstract

The major challenge for a greater dissemination of *in vitro* produced (IVP) bovine embryos is to improve embryonic survival after cryopreservation. The involvement of embryonic lipids on this issue is well documented. However, it has been recognized that not only the amount of lipids that affects embryo cryotolerance, but the embryo survival capacity after cryopreservation is a rather multifactorial event. In this review, some strategies to improve embryonic lipid composition and postcryopreservation survival by modifying the embryos themselves to make them more cryopreservable are overviewed. The use of semi-defined and defined serum-free culture media, the addition of some chemicals in the culture media to modify embryo lipid composition, and the modulation of embryo cell membrane fluidity by cholesterol or unsaturated fatty acids added to the culture media and oocyte/embryo donor nutritional management with a diet enriched in polyunsaturated fatty acids, were described as alternatives for the improvement of IVP embryo survival after cryopreservation.

**Keywords:** bovine, cryotolerance, cryosurvival, *in vitro* produced embryos, lipid.

### Introduction

The global *in vitro* production of bovine embryos has increased for the 6th consecutive year in 2011. The total number of *in vitro* produced (IVP) embryos transferred worldwide was 373,836. Brazil alone was responsible for 85% of the global market of IVP embryos (Stroud, 2012). This achievement can be attributed to the high number of oocytes recovered by ovum pick-up (OPU) of zebuine breed donors, mainly represented by Nellore (*Bos taurus indicus*) animals, allowing the commercial application of *in vitro* production on large-scale programs (Pontes *et al.*, 2011).

Embryo cryopreservation is an assisted reproductive technology that allows the storage of excess embryos derived from *in vitro* production and embryo transfer programs so they can be commercialized or transferred at the most convenient time. It is considered a strategy to overcome some logistic problems associated with the transfer of large numbers of fresh embryos and mainly for expanding the commercialization of embryos between countries (Sudano *et al.*, 2012c).

Despite the very good results associated with the fresh IVP embryo transfer, the use of cryopreserved embryos is extremely limited. The modest results of cryopreserved IVP embryos limit their application at the field conditions as it is successfully done with the semen in the artificial insemination. Even after many advances in embryo research over the past decades, embryo cryopreservation remains one of the most challenging biotechnologies of bovine reproduction, since the cryopreservation results are still inconsistent. This fact reflects directly in the lower number of cryopreserved embryos (Fig. 1) in Brazil (3 to 7%) and worldwide (7 to 8%) over the last years (Stroud, 2010, 2012; Viana *et al.*, 2010; Viana, 2012).

The most common approach to deal with the disappointing results of cryopreservation is to vary the cryopreservation procedures by altering, for example, the concentration and type of cryoprotectants, the time and temperature of the protocol, and the addition of additives (sugars or surfactants). Despite the fact that this approach usually results in improvements, they are often limited, what has led to increasing the efforts on an embryo-focused approach by modifying the embryos themselves to make them more cryopreservable (Seidel, 2006). Therefore, the objective of this review is to present some strategies for improving postcryopreservation survival capacity through an embryo-focused approach in order to produce an embryo more resistant to the cryopreservation.

<sup>4</sup>Corresponding author: mjsudano@gmail.com; fernanda@fmvz.unesp.br

Phone/Fax: +55(14)3811-6249

Received: May 10, 2013

Accepted: July 24, 2013




Figure 1. Total number and percentage of cryopreserved *in vivo* (ET) and *in vitro* (IVP) produced bovine embryos in the world (A) and Brazil (B). Data were obtained from IETS annual statistics and data retrieval committee report ([http://www.iets.org/comm\\_data.asp](http://www.iets.org/comm_data.asp)), and Brazilian record data (Viana and Camargo, 2007; Viana, 2009; Viana *et al.*, 2010; Viana, 2012).

#### Differences between *in vitro*- and *in vivo*-produced embryos and quality control of *in vitro* production systems

Since the first success of the cryopreservation of a mouse embryo (Whittingham *et al.*, 1972), several procedures were developed to cryopreserve embryos. These methods can be basically classified in two major strategies: slow freezing and vitrification. Despite that both of them are considered cryopreservation techniques, they have important conceptual differences. In the slow freezing system, extracellular water crystallizes resulting in increased osmotic gradient that draws water from the intracellular compartment until intracellular vitrification occurs. In the vitrification system, both intra and extracellular compartment vitrify after cellular dehydration has already occurred (Saragusty and Arav, 2011).

When comparing the results of these

cryopreservation methods based on the embryo origin, i.e. IVP *versus in vivo* produced (ET) embryos, ET embryos showed similar pregnancy rates (varying from 39 to 59%) for both slow freezing and vitrification procedures (Massip, 1987; van Wagtenonk-de Leeuw *et al.*, 1997; Inaba *et al.*, 2011), while the best results for IVP embryos (varying from 52 to 100% of re-expansion and from 36 to 93% of hatching/hatched rate) were achieved using vitrification (Nedambale *et al.*, 2004; Mucci *et al.*, 2006; Yu *et al.*, 2010; Inaba *et al.*, 2011). It is largely known that IVP embryos have a greater sensitivity to the cryopreservation techniques than ET ones (Leibo and Loskutoff, 1993) based on the fact that comparison of embryos from these two origins has demonstrated that the embryos do not survive equally in the different cryopreservation methods.

There are many morphological and metabolic differences between IVP and ET embryos, such as: very



electron-dense cytoplasm, loose blastomeres, buoyant density, metabolic abnormalities (“crabtree effect” and “unquite metabolism”), gene overexpression, apoptosis rate, lipid content, and postcryopreservation survival (Fair *et al.*, 2001; Abe *et al.*, 2002; Rizos *et al.*, 2002; Corcoran *et al.*, 2006; De La Torre-Sanchez *et al.*, 2006b; Mucci *et al.*, 2006; Leese *et al.*, 2008a; Côté *et al.*, 2011; Sudano *et al.*, 2011). All these alterations observed in IVP embryos can be attributed to the different *in vitro* culture conditions during oocyte maturation and embryo development that modulate the occurrence of these distinctive phenotypes (Lonergan *et al.*, 2003).

In the literature, the large amount of cytoplasmic lipid droplets observed in IVP embryos has been suggested to be the major cause of reduced postcryopreservation survival (Abe *et al.*, 2002; De La Torre-Sanchez *et al.*, 2006b; Mucci *et al.*, 2006; Barceló-Fimbres and Seidel, 2007a, b). Indeed, an increased amount of lipid droplets had a moderate correlation with the postcryopreservation survival. However, the embryo quality evaluated by the apoptosis rate had a strong correlation with the embryo survival after cryopreservation (Sudano *et al.*, 2012b), highlighting the importance of embryo quality after cryopreservation and suggesting that embryo cryosurvival capacity is a multifactorial event. Several factors are involved in the embryo cryotolerance, such as: lipid content, lipid composition, embryo metabolism, apoptosis, and global gene expression pattern (Sudano *et al.*, 2011, 2012a, b, c).

The goal during *in vitro* embryo production is try to mimick as much as possible the *in vivo* environmental condition to achieve a good quality embryo, which, in turn, could be cryopreserved more efficiently. Several researchers suggest a rigorous quality control during all steps of *in vitro* embryo production to obtain good results (Lane *et al.*, 2008; Hasler, 2010; Saragusty and Arav, 2011).

In this context, we explore in this review the important aspects that could affect IVP embryo quality and cryotolerance, namely: culture media composition (additives, salts, aminoacids, hormones, sugars, antioxidants, pH, and osmolarity), atmosphere (lower or higher oxygen tension), temperature, oocyte donor, semen, sire, and technician (Gardner, 2008; Leese *et al.*, 2008b; Feugang *et al.*, 2009; Hasler, 2010; Hugentobler *et al.*, 2010).

### Embryo lipids

There are evidences that at least four classes of lipids affect embryo survival after cryopreservation: triacylglycerides (TAG; mainly stored at the cytoplasmic lipid droplets), free fatty acids (FFA), cholesterol (Chol) and phospholipids (PL; cell membrane lipids).

The reason for the increased number of cytoplasmic lipid droplets in IVP embryos is unknown. However, it is speculated that it is related to fetal calf

serum (FCS) supplementation in the culture media. It seems that FCS increases embryo lipid content through: a) the lipoproteins from the serum are absorbed by the embryonic cells (Sata *et al.*, 1999); b) the embryo is induced to perform neosynthesis of triacylglycerides due to the presence of FCS (Razek *et al.*, 2000); and c) the FCS changes the function of  $\beta$ -oxidation in the mitochondria (Abe *et al.*, 2002). Another potential reason is that lipid accumulation occurs as an effect of abnormal energetic metabolism. An imbalance in the cellular oxidation-reduction process also occurs, affecting mitochondrial function and impairing metabolism of lipid complexes through  $\beta$ -oxidation (Abe *et al.*, 2002).

The lipid droplets present in the IVP embryos cytoplasm are mainly composed by TAG, the predominant lipid in the cytoplasm of mammalian cells (McKeegan and Sturmey, 2011). These stored lipids constitute an important source of energy for oocytes and embryos (Sturmey *et al.*, 2009). The estimated TAG content remained constant in a serum-free medium during embryo development (33 ng/embryo); however, in a serum-supplemented medium the TAG amount increased from 33 ng in 5-8 cell stage to 62 ng in hatched blastocysts (Ferguson and Leese, 1999). In addition, FCS also increased the total fatty acid amount compared with a serum-free media (74.2 vs. 57.2 ng, respectively), mostly represented by an increase in the palmitic (28.9 vs. 20.1 ng), stearic (18.0 vs. 13.1 ng), oleic (12.1 vs. 4 %), and palmitoleic (16.3 vs. 3.7 %), which are saturated (Reis *et al.*, 2003) and monounsaturated (Sata *et al.*, 1999) fatty acids.

On the other hand, PL are the most abundant lipid in eukaryotic cell membranes and their role in successful embryo cryopreservation remains poorly understood (van Meer *et al.*, 2008). Phospholipids, particularly phosphatidylcholines (PC) and sphingomyelins (SM), are structural units of functional membranes, and their composition determines the physicochemical properties of cell membranes, including fluidity, permeability, and thermal phase behavior (Edidin, 2003). We have recently reported that the PL profiles of bovine embryos vary between subspecies (*B. taurus indicus* vs. *B. taurus taurus*) and origin (IVP vs. ET) and that specific lipid species can potentially be used as biomarkers of embryonic postcryopreservation survival. These results indicate that not only the lipid amount but also the lipid composition accounts for embryo survival after cryopreservation (Sudano *et al.*, 2012c).

### Strategies to improve embryo lipid composition and postcryopreservation survival capacity

#### *Fetal calf serum-free media*

It is well known that FCS provides energy substrates, amino acids, vitamins, growth factors, and heavy-metal chelators. Although FCS has useful properties, its use has been associated with several

abnormalities, such as cell organelles modification, mitochondrial degeneration, gene expression modification, large offspring syndrome, increased lipid droplets number and reduced postcryopreservation survival (Abe *et al.*, 2002; Lazzari *et al.*, 2002; Rizos *et al.*, 2002, 2003; Sudano *et al.*, 2011). As a result, chemically-defined media (without FCS) have been developed (Keskintepe and Brackett, 1996).

It has already been described that it is possible to produce *in vitro* bovine embryos in defined (Block *et al.* 2010; Momozawa and Fukuda, 2011) or semidefined (Mucci *et al.*, 2006) serum-free media without affecting blastocyst yield and increasing embryo cryosurvival. In addition, a reduction of FCS concentration in the culture media alone was enough to decrease the lipid content and increase the postcryopreservation survival (Sudano *et al.*, 2011). The use of serum-free media has been considered as one of the first actions for the establishment of a successful *in vitro* embryo production system, allowing higher embryo survival after cryopreservation (Rizos *et al.*, 2003; Mucci *et al.*, 2006).

#### Use of chemical additives

Energetic substrate is promptly metabolized

through the glycolytic pathway (Fig. 2). However, the energetic metabolism during early embryo development (pre and post-compaction stage) is abnormal in IVP embryos (De La Torre-Sanchez *et al.*, 2006b). Under *in vitro* conditions, embryos show an increased activity of the glycolytic pathway and a consequently inhibition of oxidative phosphorylation pathway, characterized as “Crabtree effect” (Crabtree, 1929; Seshagiri and Bavister, 1991). A higher metabolic activity through the glycolysis impairs embryo development, because too little energetic substrate is partitioned to the pentose phosphate pathway (PPP) which is part of an important biosynthetic pathway (Wales and Hunter, 1990), by favoring lipid accumulation and rising cellular concentrations of lipid synthesis precursors (Rieger, 1992).

An interesting approach would be the use of phenazine ethosulfate (PES) in order to balance the energetic metabolism and reduce the lipid accumulation, by favoring the enzymatic reactions of PPP (Fig. 2), since this chemical oxidizes NADPH to NADP (De La Torre-Sanchez *et al.*, 2006a; Sudano *et al.*, 2011). The use of PES in the post-compaction period resulted in a reduction of the embryo lipid accumulation and an increase in the postcryopreservation survival (Barceló-Fimbres and Seidel, 2007b; Sudano *et al.*, 2011).




Figure 2. Energetic metabolism and phenazine ethosulfate (PES) mechanism of action in the preimplantation embryo. Energetic substrate is promptly metabolized in the glycolytic pathway (represented in black) through glycolysis within cytoplasm, followed by the krebs cycle and oxidative phosphorylation within mitochondria to produce energy in the form of ATP. PES reduces NADPH to NADP (dashed red arrows) favoring the pentose phosphate pathway (represented in green) with inhibition of the fatty acids pathway (represented in blue). Adapted from Barceló-Fimbres and Seidel (2007a).



Another chemical used to reduce lipid content of IVP embryos is the forskolin (Fig. 3A), a potent adenylate cyclase activator that stimulates the lipase activity through the cAMP/protein kinase pathway (Men *et al.*, 2006). Forskolin supplementation in the culture media reduced lipid content and increased the embryo survival after cryopreservation of bovine and porcine IVP embryos (Men *et al.*, 2006; Paschoal *et al.*, 2012).

More recently, L-carnitine, a small water-soluble molecule and cofactor of  $\beta$ -oxidation, was found to play an important role in the lipid metabolism (Sutton-McDowall *et al.*, 2012; Moawad *et al.*, 2013). This chemical is crucial for fatty acids (in form of acyl-CoA) translocation into the mitochondria (Fig. 3B), where they will be metabolized to acetyl-CoA through

$\beta$ -oxidation, and can be further metabolized in Krebs cycle and oxidative phosphorylation for ATP production (Sutton-McDowall *et al.*, 2012). L-carnitine also has an antioxidant activity protecting the cells from DNA damage (Abdelrazik *et al.*, 2009). Several beneficial effects of L-carnitine supplementation to the culture media have already been reported, including the improvement in the embryo development (Sutton-McDowall *et al.*, 2012), lipid metabolism and cryotolerance of bovine embryos (Takahashi *et al.*, 2012). The unique dual effects of L-carnitine enriching cellular lipid metabolism and providing antioxidative protection make it a chemical candidate for a non-invasive improvement of cryotolerance and developmental competence in IVP embryos (Takahashi *et al.*, 2012).

Figure 3. Mechanism of action of forskolin (A), a potent adenylate cyclase activator that stimulates the lipase activity through the cAMP / protein kinase (PKA) pathway, and L-carnitine (B), that favors fatty acid translocation into the mitochondria (in form of acyl-CoA) where they will be metabolized to acetyl-CoA through  $\beta$ -oxidation.

#### Cell membrane

In 1972, the structure of cell membrane was presented as the fluid mosaic model, which describes a cell membrane composed by a fluid bilayer of phospholipids oriented with the hydrophilic and hydrophobic portion to outside (extracellular and cytosolic compartments) and interior of the membrane, respectively (Singer and Nicolson, 1972).

The PL are composed by a three carbon backbone to which a phospho-head group is attached to

an end carbon of the backbone and two fatty acyl chains (fatty acids) are attached to the other carbons. Both the phospho-head groups and the fatty acids can vary in their composition and this will determine the properties of the cell membrane (Edidin, 2003). Indeed, it was reported that embryos with an increased abundance of unsaturated lipid species had greater cryosurvival (Sudano *et al.*, 2012c).

Cholesterol is another molecule present in cell membrane, and its level and the ratio between cholesterol and PL also affects the membrane fluidity (Horvath and



Seidel, 2006). Enriching the embryo cell membrane with unsaturated fatty acids and cholesterol to improve embryo cryotolerance has already been performed by two procedures, as follows: 1) membrane incorporation through its supplementation in the culture media; 2) nutritional management of oocyte/embryo donors and by offering a diet-rich with polyunsaturated fatty acids.

While the addition of cholesterol-loaded methyl- $\beta$ -cyclodextrin to the cryopreservation media had no effect on cryopreserved IVP bovine blastocysts (Pugh *et al.*, 1998), it seems to have a positive effect on vitrified oocytes as measured by an increase in the cleavage rate and number of eight-cell embryos (Horvath and Seidel, 2006), as well as an improvement in the nuclear maturation (Sprícigo *et al.*, 2012) after warming in comparison with the untreated group.

In addition, the unsaturated fatty acid supplementation (especially linoleic acid) in the culture media improved embryo cryotolerance (Hochi *et al.*, 1999; Pereira *et al.*, 2007) and reduced lipid content (Pereira *et al.*, 2007) of IVP embryos. Likewise, the oocyte/embryo donor nutritional management with a diet enriched in polyunsaturated fatty acids increased the cryosurvival of ewe oocytes (Zeron *et al.*, 2002) and porcine embryos (Kojima *et al.*, 1996).

### Conclusion

The major obstacle for a greater dissemination of the use of *in vitro* produced bovine embryos is their high sensitivity to the cryopreservation process. The involvement of the embryo lipids on this aspect is well documented. However, it has been recognized that not only the amount of cytoplasmic lipids affects embryo cryotolerance. The embryo survival capacity after cryopreservation is a multifactorial event. A rigorous quality control during all steps of *in vitro* embryo production is required to obtain a good quality and cryopreservable embryo. The use of a serum-free media, the addition of chemicals to change lipid metabolism, and the modulation of membrane lipid composition have been described as some alternatives for the improvement of the IVP embryo survival after cryopreservation

### Acknowledgments

The authors would like to thank Fundação de Amparo à Pesquisa do Estado de São Paulo (São Paulo Research Foundation - FAPESP), for the financial support (grants 2009/54513-3, 2010/09922-0 and 2012/02333-4).

### References

**Abdelrazik H, Sharma R, Mahfouz R, Agarwal A.** 2009. L-carnitine decreases DNA damage and improves the *in vitro* blastocyst development rate in mouse embryos. *Fertil Steril*, 91:589-596.

**Abe H, Yamashita S, Satoh T, Hoshi H.** 2002.

Accumulation of cytoplasmic lipid droplets in bovine embryos and cryotolerance of embryos developed in different culture systems using serum-free or serum-containing media. *Mol Reprod Dev*, 61:57-66.

**Barceló-Fimbres M, Seidel GE.** 2007a. Effects of either glucose or fructose and metabolic regulators on bovine embryo development and lipid accumulation *in vitro*. *Mol Reprod Dev*, 74:1406-1418.

**Barceló-Fimbres M, Seidel GE.** 2007b. Effects of fetal calf serum, phenazine ethosulfate and either glucose or fructose during *in vitro* culture of bovine embryos on embryonic development after cryopreservation. *Mol Reprod Dev*, 74:1395-1405.

**Block J, Bonilla L, Hansen PJ.** 2010. Efficacy of *in vitro* embryo transfer in lactating dairy cows using fresh or vitrified embryos produced in a novel embryo culture medium. *J Dairy Sci*, 93:5234-5242.

**Corcoran D, Fair T, Park S, Rizos D, Patel OV, Smith GW, Coussens PM, Ireland JJ, Boland MP, Evans AC, Lonergan P.** 2006. Suppressed expression of genes involved in transcription and translation in *in vitro* compared with *in vivo* cultured bovine embryos. *Reproduction*, 131:651-660.

**Crabtree HG.** 1929. Observations on the carbohydrate metabolism of tumours. *Biochem J*, 23:536-545.

**Côté I, Vigneault C, Laflamme I, Laquerre J, Fournier É, Gilbert I, Scantland S, Gagné D, Blondin P, Robert C.** 2011. Comprehensive cross production system assessment of the impact of *in vitro* microenvironment on the expression of messengers and long non-coding RNAs in the bovine blastocyst. *Reproduction*, 142:99-112.

**De La Torre-Sanchez JF, Gardner DK, Preis K, Gibbons J, Seidel GE.** 2006a. Metabolic regulation of *in vitro*-produced bovine embryos. II. Effects of phenazine ethosulfate, sodium azide and 2,4-dinitrophenol during post-compaction development on glucose metabolism and lipid accumulation. *Reprod Fertil Dev*, 18:597-607.

**De La Torre-Sanchez JF, Preis K, Seidel GE.** 2006b. Metabolic regulation of *in vitro*-produced bovine embryos. I. Effects of metabolic regulators at different glucose concentrations with embryos produced by semen from different bulls. *Reprod Fertil Dev*, 18:585-596.

**Edidin M.** 2003. Lipids on the frontier: a century of cell-membrane bilayers. *Nat Rev Mol Cell Biol*, 4:414-418.

**Fair T, Lonergan P, Dinnyes A, Cottell DC, Hyttel P, Ward FA, Boland MP.** 2001. Ultrastructure of bovine blastocysts following cryopreservation: effect of method of blastocyst production. *Mol Reprod Dev*, 58:186-195.

**Ferguson EM, Leese HJ.** 1999. Triglyceride content of bovine oocytes and early embryos. *J Reprod Fertil*, 116:373-378.

**Feugang JM, Camargo-Rodriguez O, Memili E.** 2009. Culture systems for bovine embryos. *Livest Sci*, 121:141-149.

**Gardner DK.** 2008. Dissection of culture media for embryos: the most important and less important



- components and characteristics. *Reprod Fertil Dev*, 20:9-18.
- Hasler JF.** 2010. Synthetic media for culture, freezing and vitrification of bovine embryos. *Reprod Fertil Dev*, 22:119-125.
- Hochi S, Kimura K, Hanada A.** 1999. Effect of linoleic acid-albumin in the culture medium on freezing sensitivity of in vitro-produced bovine morulae. *Theriogenology*, 52:497-504.
- Horvath G, Seidel GE.** 2006. Vitrification of bovine oocytes after treatment with cholesterol-loaded methyl-beta-cyclodextrin. *Theriogenology*, 66:1026-1033.
- Hugentobler SA, Sreenan JM, Humpherson PG, Leese HJ, Diskin MG, Morris DG.** 2010. Effects of changes in the concentration of systemic progesterone on ions, amino acids and energy substrates in cattle oviduct and uterine fluid and blood. *Reprod Fertil Dev*, 22:684-694.
- Inaba Y, Aikawa Y, Hirai T, Hashiyada Y, Yamanouchi T, Misumi K, Ohtake M, Somfai T, Kobayashi S, Saito N, Matoba S, Konishi K, Imai K.** 2011. In-straw cryoprotectant dilution for bovine embryos vitrified using Cryotop. *J Reprod Dev*, 57:437-443.
- Keskinetepe L, Brackett BG.** 1996. In vitro developmental competence of in vitro-matured bovine oocytes fertilized and cultured in completely defined media. *Biol Reprod*, 55:333-339.
- Kojima T, Iwasaki T, Zeniya Y, Yoshino J, Totsukawa K.** 1996. Effect of dietary administration with linoleic acid and alpha-tocopherol on freezing tolerance in porcine embryos. *J Reprod Dev*, 42:67-72.
- Lane M, Mitchell M, Cashman KS, Feil D, Wakefield S, Zander-Fox DL.** 2008. To QC or not to QC: the key to a consistent laboratory? *Reprod Fertil Dev*, 20:23-32.
- Lazzari G, Wrenzycki C, Herrmann D, Duchi R, Kruip T, Niemann H, Galli C.** 2002. Cellular and molecular deviations in bovine in vitro-produced embryos are related to the large offspring syndrome. *Biol Reprod*, 67:767-775.
- Leese HJ, Baumann CG, Brison DR, McEvoy TG, Sturmey RG.** 2008a. Metabolism of the viable mammalian embryo: quietness revisited. *Mol Hum Reprod*, 14:667-672.
- Leese HJ, Hugentobler SA, Gray SM, Morris DG, Sturmey RG, Whitear SL, Sreenan JM.** 2008b. Female reproductive tract fluids: composition, mechanism of formation and potential role in the developmental origins of health and disease. *Reprod Fertil Dev*, 20:1-8.
- Leibo SP, Loskutoff NM.** 1993. Cryobiology of in vitro derived bovine embryos. *Theriogenology*, 39:81-94.
- Lonergan P, Rizos D, Kanka J, Nemcova L, Mbaye AM, Kingston M, Wade M, Duffy P, Boland MP.** 2003. Temporal sensitivity of bovine embryos to culture environment after fertilization and the implications for blastocyst quality. *Reproduction*, 126:337-346.
- Massip A.** 1987. Recent progress in cryopreservation of cattle embryos. *Theriogenology*, 27:69-79.
- McKeegan PJ, Sturmey RG.** 2011. The role of fatty acids in oocyte and early embryo development. *Reprod Fertil Dev*, 24:59-67.
- Men H, Agca Y, Riley LK, Critser JK.** 2006. Improved survival of vitrified porcine embryos after partial delipitation through chemically stimulated lipolysis and inhibition of apoptosis. *Theriogenology*, 66:2008-2016.
- Moawad AR, Tan SL, Xu B, Chen HY, Taketo T.** 2013. L-carnitine supplementation during vitrification of mouse oocytes at the germinal vesicle stage improves pre-implantation development following maturation and fertilization in vitro. *Biol Reprod*, 88:104.
- Momozawa K, Fukuda Y.** 2011. Establishment of an advanced chemically defined medium for early embryos derived from in vitro matured and fertilized bovine oocytes. *J Reprod Dev*, 57:681-689.
- Mucci N, Aller J, Kaiser GG, Hozbor F, Cabodevila J, Alberio RH.** 2006. Effect of estrous cow serum during bovine embryo culture on blastocyst development and cryotolerance after slow freezing or vitrification. *Theriogenology*, 65:1551-1562.
- Nedambale TL, Dinnyés A, Groen W, Dobrinsky JR, Tian XC, Yang X.** 2004. Comparison on in vitro fertilized bovine embryos cultured in KSOM or SOF and cryopreserved by slow freezing or vitrification. *Theriogenology*, 62:437-449.
- Paschoal DM, Sudano MJ, Guastali MD, Dias Maziero RR, Crocomo LF, Oña Magalhães LC, Silva Rascado T, Martins A, Landim-Alvarenga FD.** 2012. Forskolin effect on the cryosurvival of in vitro-produced bovine embryos in the presence or absence of fetal calf serum. *Zygote*, 18:1-12.
- Pereira RM, Baptista MC, Vasques MI, Horta AE, Portugal PV, Bessa RJ, Silva JC, Pereira MS, Marques CC.** 2007. Cryosurvival of bovine blastocysts is enhanced by culture with trans-10 cis-12 conjugated linoleic acid (10t,12c CLA). *Anim Reprod Sci*, 98:293-301.
- Pontes JH, Melo Sterza FA, Basso AC, Ferreira CR, Sanches BV, Rubin KC, Seneda MM.** 2011. Ovum pick up, in vitro embryo production, and pregnancy rates from a large-scale commercial program using Nelore cattle (*Bos indicus*) donors. *Theriogenology*, 75:1640-1646.
- Pugh PA, Ankersmit AE, McGowan LT, Tervit HR.** 1998. Cryopreservation of in vitro-produced bovine embryos: effects of protein type and concentration during freezing or of liposomes during culture on post-thaw survival. *Theriogenology*, 50:495-506.
- Razek IAE, Charpigny G, Kodja S, Marquant-Leguienne B, Mermillod P, Guyader J, Oly C.** 2000. Differences in lipid composition between in vivo- and in vitro-produced bovine embryos. *Theriogenology*, 53:346.
- Reis A, Rooke JA, McCallum GJ, Staines ME, Ewen**



- M, Lomax MA, McEvoy TG.** 2003. Consequences of exposure to serum, with or without vitamin E supplementation, in terms of the fatty acid content and viability of bovine blastocysts produced in vitro. *Reprod Fertil Dev*, 15:275-284.
- Rieger D.** 1992. Relationships between energy metabolism and development of early mammalian embryos. *Theriogenology*, 37:75-93.
- Rizos D, Ward F, Duffy P, Boland MP, Lonergan P.** 2002. Consequences of bovine oocyte maturation, fertilization or early embryo development in vitro versus in vivo: implications for blastocyst yield and blastocyst quality. *Mol Reprod Dev*, 61:234-248.
- Rizos D, Gutiérrez-Adán A, Pérez-Garnelo S, De La Fuente J, Boland MP, Lonergan P.** 2003. Bovine embryo culture in the presence or absence of serum: implications for blastocyst development, cryotolerance, and messenger RNA expression. *Biol Reprod*, 68:236-243.
- Saragusty J, Arav A.** 2011. Current progress in oocyte and embryo cryopreservation by slow freezing and vitrification. *Reproduction*, 141:1-19.
- Sata R, Tsujii H, Abe H, Yamashita S, Hoshi H.** 1999. Fatty acid composition of bovine embryos cultured in serum-free and serum containing medium during early embryonic development. *J Reprod Dev*, 45:97-103.
- Seidel GE.** 2006. Modifying oocytes and embryos to improve their cryopreservation. *Theriogenology*, 65:228-235.
- Seshagiri PB, Bavister BD.** 1991. Glucose and phosphate inhibit respiration and oxidative metabolism in cultured hamster eight-cell embryos: evidence for the "crabtree effect". *Mol Reprod Dev*, 30:105-111.
- Singer SJ, Nicolson GL.** 1972. The fluid mosaic model of the structure of cell membranes. *Science*, 175:720-731.
- Sprícigo JF, Morais KS, Yang BS, Dode MA.** 2012. Effect of the exposure to methyl- $\beta$ -cyclodextrin prior to chilling or vitrification on the viability of bovine immature oocytes. *Cryobiology*, 65:319-325.
- Stroud B.** 2010. IETS Statistics and Data Retrieval Committee Report. Champaign, IL: IETS.
- Stroud B.** 2012. IETS 2012 Statistics and Data Retrieval Committee Report. Champaign, IL: IETS.
- Sturmev RG, Reis A, Leese HJ, McEvoy TG.** 2009. Role of fatty acids in energy provision during oocyte maturation and early embryo development. *Reprod Domest Anim*, 44(suppl. 3):50-58.
- Sudano MJ, Paschoal DM, Rascado TD, Magalhaes LCO, Crocomo LF, Lima-Neto JF, Landim-Alvarenga FD.** 2011. Lipid content and apoptosis of in vitro-produced bovine embryos as determinants of susceptibility to vitrification. *Theriogenology*, 75:1211-1220.
- Sudano MJ, Paschoal DM, Caixeta ES, Santos VG, Tata A, Ferreira CR, Martins Junior A, Machado R, Eberlin MN, Buratini J, Landim-Alvarenga FC.** 2012a. Cryotolerance of *Bos taurus indicus* and *Bos taurus taurus* in vitro and in vivo produced embryos. *Anim Reprod*, 9:677. (abstract).
- Sudano MJ, Paschoal DM, Silva Rascado T, Crocomo LF, Magalhães LC, Martins Junior A, Machado R, Landim-Alvarenga FD.** 2012b. Crucial surviving aspects for vitrified in vitro-produced bovine embryos. *Zygote*, 11:1-8.
- Sudano MJ, Santos VG, Tata A, Ferreira CR, Paschoal DM, Machado R, Buratini J, Eberlin MN, Landim-Alvarenga FD.** 2012c. Phosphatidylcholine and sphingomyelin profiles vary in *Bos taurus indicus* and *Bos taurus taurus* in vitro- and in vivo-produced blastocysts. *Biol Reprod*, 87:130.
- Sutton-McDowall ML, Feil D, Robker RL, Thompson JG, Dunning KR.** 2012. Utilization of endogenous fatty acid stores for energy production in bovine preimplantation embryos. *Theriogenology*, 77:1632-1641.
- Takahashi T, Inaba Y, Somfai T, Kaneda M, Geshi M, Nagai T, Manabe N.** 2012. Supplementation of culture medium with l-carnitine improves development and cryotolerance of bovine embryos produced in vitro. *Reprod Fertil Dev*, 25:589-599.
- van Meer G, Voelker DR, Feigenson GW.** 2008. Membrane lipids: where they are and how they behave. *Nat Rev Mol Cell Biol*, 9:112-124.
- van Wagtenonk-de Leeuw AM, den Daas JH, Rall WF.** 1997. Field trial to compare pregnancy rates of bovine embryo cryopreservation methods: vitrification and one-step dilution versus slow freezing and three-step dilution. *Theriogenology*, 48:1071-1084.
- Viana JHM, Camargo LSA.** 2007. Bovine embryo production in Brazil: a new scenario. *Acta Sci Vet*, 35(suppl):s915-s924.
- Viana JHM.** 2009. Mudanças e tendências no mercado de embriões bovinos no Brasil. *Embrião*, 42:5-7.
- Viana JHV, Siqueira LGB, Palhão MP, Camargo LSA.** 2010. Use of in vitro fertilization technique in the last decade and its effect on brazilian embryo industry and animal production. *Acta Sci Vet*, 38(suppl):s661-s674.
- Viana JHV.** 2012. Levantamento estatístico da produção de embriões bovinos no Brasil em 2011: mudanças e tendências futuras. *Embrião*, 51:6-10.
- Wales RG, Hunter J.** 1990. Participation of glucose in the synthesis of glycoproteins in preimplantation mouse embryos. *Reprod Fertil Dev*, 2:35-50.
- Whittingham DG, Leibo SP, Mazur P.** 1972. Survival of mouse embryos frozen to -196 degrees and -269 degrees C. *Science*, 178:411-414.
- Yu XL, Deng W, Liu FJ, Li YH, Li XX, Zhang YL, Zan LS.** 2010. Closed pulled straw vitrification of in vitro-produced and in vivo-produced bovine embryos. *Theriogenology*, 73:474-479.
- Zeron Y, Sklan D, Arav A.** 2002. Effect of polyunsaturated fatty acid supplementation on biophysical parameters and chilling sensitivity of ewe oocytes. *Mol Reprod Dev*, 61:271-278.