



## Fertility programs for lactating dairy cows, their physiological basis, and the factors that are critical for their success

J.P.N. Martins, J.R. Pursley<sup>1</sup>

Department of Animal Science, Michigan State University, East Lansing, MI 48824.

### Abstract

Lactating dairy cows have unique reproductive responses compared to when they were heifers that result in distinctly different reproductive measurements and pregnancy outcomes that can be partially overcome with pharmacological strategies. These parameters include circulating progesterone and estradiol concentrations, ovulatory follicle and corpus luteum diameter, incidence of anovulation and double ovulations, time in estrus, pregnancies per artificial insemination and pregnancy losses. Circulating concentrations of progesterone during diestrus are approximately half that in cows compared with heifers. This marked difference in progesterone is likely the explanation for an increased size in diameter of the ovulatory follicle and incidence of double ovulations in cows compared with heifers. Differences in diameter of the ovulatory follicle may explain why cows have greater corpora lutea diameters compared to heifers. The increase in double ovulations appears to be a key driver in the increase in twinning and pregnancy loss as dairy heifers transition to primi- and multiparous cows. Reduced estradiol concentrations in cows at time of estrus helps to explain the decreased duration of estrus in cows compared with heifers. Concentrations of progesterone during growth of the ovulatory follicle may be a key driver in differences in pregnancies per AI in cows compared with heifers. The difference in circulating progesterone may be related to LH overstimulation of the oocyte/cumulus complex in cows compared to when they were heifers. Pharmacological strategies have been developed in lactating dairy cows to manipulate ovarian development to create a hormonal environment similar to that of heifers. Three primary strategies, Presynch-11, G6G and Double Ovsynch appear to enhance fertility of dairy cows. This review discusses how these three strategies manipulate ovarian development similar to that of heifers and why compelling data indicate these programs should be referred to as “fertility programs.”

**Keywords:** corpus luteum, dairy, follicle, ovsynch, progesterone.

### Introduction

The evolutionary role of the ovary is to produce oocytes capable of fertilization. The supporting cast of ovarian structures, follicles and corpora lutea, are responsible for the hormonal environment, production of oocytes and maintenance of pregnancy. The physiological outcomes of these structures in lactating

dairy cows when compared with nulliparous heifers is associated with negative reproductive outcomes (Wiltbank *et al.*, 2006). Pharmacologic manipulation of ovarian structures can reverse these effects of lactation and enhance fertility (Wiltbank *et al.*, 2011). The physiological basis for development of pharmacological programs to improve fertility of lactating dairy cows comes from studies that characterized key differences in reproductive indices in primiparous and multiparous cows (referred heretofore as “cows”) vs. nulliparous heifers (referred heretofore as “heifers”; Sartori *et al.*, 2002, 2004; Wolfenson *et al.*, 2004). Differences in circulating concentrations of progesterone (P4) appears to be the key driver of many of these indices. Lactating cows have reduced circulating concentrations of P4 during the estrous cycle compared to heifers (Sartori *et al.*, 2002, 2004; Wolfenson *et al.*, 2004). This likely results in greater number of pulses of LH during the luteal phase of the cycle and in turn drives the growth of larger dominant and ovulatory follicle diameters in cows compared with heifers (Bergfeld *et al.*, 1996). These differences in P4 and LH pulsatility also create differences in length of follicular waves (Wolfenson *et al.*, 2004). In this case, cows with lower concentrations of P4 have longer inter-wave intervals compared to heifers (Wolfenson *et al.*, 2004) due in part to increased numbers of LH pulses that drive the growth of a dominant follicle for longer periods. This, in turn, leads to more cows with an ovulatory follicle with a greater antral age (measured from onset of wave to ovulation) that developed under greater numbers of LH pulses compared to heifers. Heifers have greater chances to have three waves of follicle growth during a slightly shorter estrous cycle if the second wave dominant follicle becomes atretic prior to endogenous luteolysis (Savio *et al.*, 1990). An increase in double ovulations as dairy heifers transition to primi- and multiparous cow appears to be a key driver in the increase in twinning (Wiltbank *et al.*, 2000, 2006; Sartori *et al.*, 2004; Lopez *et al.*, 2005). It appears the increase in twinning has a significant impact on pregnancy loss (López-Gatius *et al.*, 2002; López-Gatius and Hunter, 2005). Metabolic differences such as milk production and increased dry matter intake may explain these differences in reproductive function (Wiltbank *et al.*, 2006).

Reproductive inefficiency is an obstacle to dairy farm profitability and sustainability. During the past 50 years, reproductive efficiency of lactating dairy cows progressively decreased due primarily to two key reproductive parameters, low estrus detection and pregnancies per AI (Lucy, 2001; Washburn *et al.*, 2002). Current reports indicate that estrus detection rate

<sup>1</sup>Corresponding author: pursleyr@msu.edu

Received: July 5, 2016

Accepted: July 28, 2016



is ~36% in lactating cows and 70% in heifers. The dramatic change in estrus detection rate from the transition of heifer to cow may be attributed to changes in circulating concentrations of estrogen (Sartori *et al.*, 2004; Wolfenson *et al.*, 2004; Wiltbank *et al.*, 2006) as well as differences in environment, with cows spending more time on concrete (Platz *et al.*, 2008; Martin *et al.*, 2015) and being more susceptible to heat stress (Her *et al.*, 1988; Orihuela, 2000). Fortunately, estrus detection rate (or service rate when referring to cows timed-inseminated) is significantly enhanced when cows are timed-inseminated with Ovsynch (Pursley *et al.*, 1997a). This is because Ovsynch can be utilized as a tool to control time to first AI and subsequent inseminations following a negative pregnancy diagnosis. Unfortunately, the second greatest obstacle, pregnancies per AI, is not enhanced with Ovsynch (Pursley *et al.*, 1997a, b). In the past 40 years, pregnancies per AI in lactating dairy cows decreased from around 65% in the 1950's (Spalding *et al.*, 1975; Butler and Smith, 1989) to approximately 35% (Strickland *et al.*, 2010; Fricke *et al.*, 2014) while pregnancies per AI in heifers remained steady at about 70% (Pursley *et al.*, 1997b; Escalante *et al.*, 2013). Ovsynch was developed to synchronize the time of ovulation to allow for timed-AI. Now, studies have focused on ways to improve follicular and luteal dynamics during Ovsynch to improve pregnancies per AI of lactating dairy cows (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006; Wiltbank and Pursley, 2014).

Three key physiological events transpire in fertility programs in a greater percentage of cows than in conventional synchronization programs: 1) A new estrous cycle and 2) subsequent ovulation of a first wave dominant follicle are induced within an 8-h period. And, 3) complete luteolysis is controlled prior to endogenously induced luteolysis. These three events must occur to make these programs different than other synchronization programs.

This review focuses on specific pharmacological interventions utilizing only gonadotropin-releasing hormone (GnRH) and prostaglandin F<sub>2</sub> $\alpha$  (PGF<sub>2</sub> $\alpha$ ) to manipulate ovarian development in lactating cows to generate physiological outcomes similar to nulliparous heifers and enhance fertility. These interventions manipulate antral age of the ovulatory follicle and the hormonal environment during its growth necessary to induce ovulation of a single competent oocyte. The outcomes improve the chances of a pregnancy to a single AI compared to AI following estrus and are now referred to as Ovsynch-based "fertility programs for dairy cows." (Wiltbank and Pursley, 2014) They are distinguishable from other Ovsynch-based synchronization programs in the way follicles and corpora lutea are manipulated.

### Limitations of solely using Ovsynch

Ovsynch is based on three pharmacological treatments (Pursley *et al.*, 1995). The first treatment, GnRH, may induce an LH surge and may cause a mature functional dominant follicle(s; DF) to ovulate (Sartori *et al.*, 2001). In turn, ovulation of the DF

induces subsequent emergence of a new follicular wave ~1.5 days later (Pursley *et al.*, 1995) followed by development of a new dominant follicle during the next 7 days. If a follicle does not respond to the GnRH it is likely the cow is in the first 3 to 4 days of a follicular wave (Bello *et al.*, 2006). During this early stage of a wave, the largest growing follicle may be too immature (e.g., no LH receptors (Xu *et al.*, 1995) to respond to the GnRH-induced LH surge caused by the first GnRH treatment (Sartori *et al.*, 2001). If the DF is not responsive to a GnRH induced LH surge, it may develop into an ovulatory follicle during the remainder of the Ovsynch treatments or possibly become atretic prior to luteolysis (Vasconcelos *et al.*, 2003). If the follicle becomes atretic prior to PGF<sub>2</sub> $\alpha$ , a new follicular wave will emerge. The new DF from that wave will most likely not be mature enough 2 days later to respond to the LH surge induced by the final GnRH of Ovsynch. If the PGF<sub>2</sub> $\alpha$  induces complete luteolysis, cows will likely display signs of estrus 3 to 4 days following AI following maturation of the newest pre-ovulatory follicle. This artifact is a common asynchrony of Ovsynch.

The second treatment, PGF<sub>2</sub> $\alpha$ , is administered to induce luteolysis, thus enabling the DF of the *new* follicular wave to develop into a pre-ovulatory follicle. Luteolysis following a single dose of PGF<sub>2</sub> $\alpha$  may not be effective, particularly in multiparous cows. This will be discussed in more detail later in this paper. The third treatment, GnRH, is administered 56 to 60 h after PGF<sub>2</sub> $\alpha$  (Brusveen *et al.*, 2008) to induce a pre-ovulatory LH surge that triggers ovulation of the DF 24 to 32 h later (Pursley *et al.*, 1995). This is highly effective in cows with functional pre-ovulatory follicles.

Cows treated with Ovsynch yield overall pregnancies per AI similar to those obtained after breeding to detected estrus (37 vs. 39%, respectively;  $P > 0.10$ ; Pursley *et al.*, 1997a). Up to 40% of cows may not have an ovulation synchronized with Ovsynch programs (Pursley *et al.*, 1995; Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). Non-synchronized cows will not be inseminated at an appropriate time relative to ovulation, or may not have luteolysis, thereby decreasing their chances of becoming pregnant (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). Improving synchronization rates alone of Ovsynch could have a dynamic impact on reproductive performance. Vasconcelos *et al.* (1999) attributed most of the variability in synchronization rate in cows to the stage of the estrous cycle in which Ovsynch was initiated. Cows started on Ovsynch between d 5-9 of the estrous cycle had a greater probability of synchronizing and therefore had a greater chance of pregnancy (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006).

### Difference between fertility and synchronization programs

Studies indicate that synchronization rates can be significantly improved when lactating dairy cows are treated with a pre-synchronization program utilizing PGF<sub>2</sub> $\alpha$  and GnRH compared to Ovsynch alone or



Presynch-12 or -14/Ovsynch (Bello *et al.*, 2006; Souza *et al.*, 2008; Herlihy *et al.*, 2012). The key reason for greater synchrony was the synchronous initiation of a new estrous cycle, which allowed the first GnRH of Ovsynch to be administered near day 6 or 7 of the new cycle (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). The data presented next will demonstrate that initiating the first GnRH of Ovsynch near day 6 or 7 of the estrous cycle enhances function of the ovulatory follicle and increases the percentage of cows that have ovulation to the final GnRH of Ovsynch (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). This, in turn, allowed more cows the opportunity for pregnancy, and translated into increased pregnancies per AI (Bello *et al.*, 2006).

*Control of the emergence of the ovulatory follicle is critical for optimal synchronization*

Attaining consistent ovulation in response to first GnRH of Ovsynch constitutes the first key step to optimizing synchronization of ovulation to Ovsynch in lactating dairy cows (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). Ovulation to first GnRH of Ovsynch is followed by emergence of a new follicular wave, from which the ovulatory follicle of Ovsynch develops (Pursley *et al.*, 1995). Thus, variation in response to first GnRH leads to extreme variation in the timing of emergence of the ovulatory follicle of Ovsynch. This, in turn, results in substantial variation in size of ovulatory follicles at the time of the final GnRH of Ovsynch (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). This variation leads to a reduced chance of pregnancy (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006).

The G6G program, for example, decreased variability in size of the ovulatory follicle and increased synchronization rate to Ovsynch (Bello *et al.*, 2006). In these experiments, cows were treated with 25 mg PGF2 $\alpha$  then 2 days later 100  $\mu$ g GnRH. Then, cows received the first GnRH of Ovsynch either 4, 5, or 6 days later in experiment 1 (Bello *et al.*, 2006), and either 6, 7 or 8 days later in experiment 2 (Bello and Pursley, 2007). Controls in both experiments received only Ovsynch. Compared to Ovsynch alone, 6 days from the presynchronization treatment of GnRH until the first GnRH of Ovsynch significantly improved percentage of cows ovulating to first GnRH, percentage of cows responding to PGF2 $\alpha$  by luteolysis, and percentage of cows with both a luteolytic response to PGF2 $\alpha$  and ovulation to the final GnRH of Ovsynch. These improvements were repeated in experiment 2 for day 6 compared to controls. In the two studies combined, ovulation rate to the first GnRH of Ovsynch averaged 90% in the day 6 groups (n = 76). In addition, it appears that 7 days from presynchronization GnRH to the first GnRH of Ovsynch also improved these responses, particularly when cows initiated a new estrous cycle by responding to both presynchronization treatments. Thus, day 6 or 7 of the estrous cycle appear to be the ideal d of the estrous cycle to initiate Ovsynch to maximize ovulatory response to the first GnRH and

luteolysis following PGF2 $\alpha$  (Bello *et al.*, 2006; Bello and Pursley, 2007).

Additional data reveal that cows ovulating in response to the first GnRH of Ovsynch yielded significantly less variability in pre-ovulatory follicle size at the final treatment of GnRH, a greater chance of luteolysis in response to the PGF2 $\alpha$  of Ovsynch, and a greater chance of ovulating to the final GnRH (Vasconcelos *et al.*, 1999; Bello *et al.*, 2006). Also from this study (Bello *et al.*, 2006), a positive linear relationship was detected between concentrations of estradiol at the final GnRH of Ovsynch and the probability of pregnancy. In addition, a quadratic relationship was also detected between ovulatory follicle size at final GnRH and the probability of a pregnancy. Cows with follicle sizes associated with a greater chance of pregnancy also had greater serum concentrations of estradiol. Thus, it is of critical importance to optimize the size of the ovulatory follicle to allow these follicles to secrete as much estradiol as possible at the time of the final treatment of GnRH of Ovsynch.

These two experiments were designed to only test the impact of this presynchronization scheme on follicle and CL development in response to the Ovsynch treatments (Bello *et al.*, 2006; Bello and Pursley, 2007). In these preliminary data, we show nearly a doubling of percent cows pregnant in the 6 or 7 days groups compared to Ovsynch alone. Figure 1 describes some of the potential differences between two presynchronization schemes. Presynch utilizes two injections of PGF2 $\alpha$  14 days apart and 11 to 14 days prior to the start of Ovsynch. Since PGF2 $\alpha$  only directly controls luteolysis, time to estrus and ovulation can be quite variable, as a result d of the cycle at the start of Ovsynch can be variable too. The likelihood of initiating ovulation to the first GnRH of Ovsynch is approximately 61% in the 11 days interval, and approximately 45% in the 14 days interval, between second PGF2 $\alpha$  and first GnRH of Ovsynch (Galvão *et al.*, 2007). In addition, cows treated with Ovsynch that are anovular will not respond to the PGF2 $\alpha$  injections of Presynch, but will likely respond to the GnRH of the new proposed program, thus allowing the initiation of Ovsynch at the optimal time of a subsequent follicular wave.

If cows were on day 6 of the cycle at time of first GnRH of Ovsynch, 97% of cows had accessory CL induced from the GnRH-induced LH surge, had significantly greater P4 concentrations, and had a greater probability of a pregnancy (Bello *et al.*, 2006; Bello and Pursley, 2007). Cows with both a day 7 and 13 corpora lutea at time of PGF2 $\alpha$  of Ovsynch have approximately 50% greater P4 concentrations compared to cows with only a day 13 corpus luteum (Pursley and Martins, 2011). Fertility programs enhance the percentage of cows that respond to the first GnRH of Ovsynch, and in turn, allows for more cows with accessory CL, greater concentrations of P4 at time of induced luteolysis, and a greater chance for pregnancy.




Figure 1. Description of potential difference in control of follicle development in Presynch versus the PGF2 $\alpha$  – GnRH presynchronization scheme proposed. Control of a new estrous cycle in more cows may allow the PGF2 $\alpha$  – GnRH scheme greater control of the ovulatory follicle following the first GnRH of Ovsynch. PG: PGF2 $\alpha$ ; F: Dominant follicle; AI: artificial insemination.

### Enhancing CL regression

Circulating concentration of P4 during fertility treatments appears to be one of the most important markers of subsequent pregnancy success following timed-AI. In addition to the effect of levels of P4 prior to PGF2 $\alpha$  on fertility of lactating dairy cows previously discussed; serum concentrations of P4 following PGF2 $\alpha$  of Ovsynch has also been associated with P/AI of lactating dairy cows after timed-AI (Souza *et al.*, 2007; Brusveen *et al.*, 2009; Martins *et al.*, 2011a). However, in this case, a slight increase on serum concentration of P4 appears to be detrimental for the success of timed-AI outcomes. Studies using synchronization programs reported that cows with functional CL that do not decrease circulating P4 to basal levels have small to no chances of conceiving following timed-AI. In previous studies, probability of pregnancy decreased as circulating concentrations of P4 increased (Souza *et al.*, 2007; Brusveen *et al.*, 2009; Martins *et al.*, 2011a). Time to reach complete luteolysis after PGF2 $\alpha$  of Ovsynch appears to also influence fertility since cows with a delay on P4 clearance had impaired fertility following timed-AI (Martins *et al.*, 2011a, b). Therefore, it is essential that cows with functional CL

have complete luteal regression following PGF2 $\alpha$  of Ovsynch, which is characterized as a drop of circulating P4 to basal levels prior to timed-AI. Reports of percentage of lactating dairy cows without complete luteolysis following PGF2 $\alpha$  of Ovsynch are between 5 and 30% (Souza *et al.*, 2007; Brusveen *et al.*, 2009; Bisinotto *et al.*, 2010; Martins *et al.*, 2011a; Giordano *et al.*, 2012). In addition, similar results were obtained using either of the two PGF2 $\alpha$  products available in the U.S.: dinoprost tromethamine (Lutalyse and ProstaMate) and cloprostenol sodium (Estrumate and estroPLAN). Although the mechanisms involved on the resistance of a mature day 7 or older CL to undergo complete luteolysis are not well characterized, some factors appear to influence the proportion of cows with complete luteolysis following PGF2 $\alpha$  of Ovsynch. A previous study from our laboratory identified that parity and service could affect percentage of cows with complete luteolysis (Martins *et al.*, 2011a). A greater percentage of cows in first service underwent luteolysis compared to second and greater services (79 vs. 71%, respectively). Primiparous cows were also more likely to have complete luteolysis compared to multiparous cows (94 vs. 81%; Martins *et al.*, 2011a). This same study also reported that cows with greater circulating of



P4 at time of PGF2 $\alpha$  of Ovsynch had a greater probability of complete luteal regression (Fig. 2; Martins *et al.*, 2011a). This result was unexpected since cows with two CL, a mature and accessory CL, at time of PGF2 $\alpha$  of Ovsynch have higher serum P4 at time of PGF2 $\alpha$  of Ovsynch and were believed to have problems with luteolysis due the number of CL and the young age of the accessory CL (day 7).

In order to enhance percentage of cows with complete luteolysis following PGF2 $\alpha$  of Ovsynch, two different approaches have been tested: repeated administrations with PGF2 $\alpha$  commercial label dose (Brusveen *et al.*, 2009) and increased label dose of PGF2 $\alpha$  administered once (Giordano *et al.*, 2013). Percentage of cows with complete CL regression was increased when an additional PGF2 $\alpha$  treatment was administered 24 h after the PGF2 $\alpha$  of Ovsynch (95.6%) compared to cows with only the PGF2 $\alpha$  of Ovsynch (84.6%; Brusveen *et al.*, 2009). There was an increase in percentage of cows with luteolysis. P/AI in cows with two vs. one PGF2 $\alpha$  treatment was 52.7 vs. 47.0% (Brusveen *et al.*, 2009). Based on data from Martins *et al.* (2011a), cows that do not have complete luteolysis have nearly a 0% chance of becoming pregnant. The 11% of cows that did not have complete

luteolysis (95.6 - 84.6%) likely did not become pregnant and mathematically this calculates to nearly a 6% difference.

Giordano *et al.* (2013) tested if a 50% increase in the label dose (0.5 mg vs. 0.75 mg) of a PGF2 $\alpha$  analogue (cloprostenol) would increase the percentage of cows with complete luteolysis following PGF2 $\alpha$  of Ovsynch (Giordano *et al.*, 2013). The greater dose of PGF2 $\alpha$  increased the percentage of cows with complete luteolysis (87.7 vs. 79.2%) and tended to increase P/AI 39 d after AI in multiparous cows (45.4 vs. 40.9%; Giordano *et al.*, 2013). However, it did not influence primiparous cows (92.8 vs. 89.7%; Giordano *et al.*, 2013). These studies indicated that some cows and/or their CL are more resistant to achieve complete luteolysis with the regular label dose of PGF2 $\alpha$ . Taken together, there is compelling evidence indicating that insufficient luteolysis after PGF2 $\alpha$  of Ovsynch has a direct impact on reproductive performance of lactating dairy cow and that two administrations of PGF2 $\alpha$  24 h apart can overcome this problem. Therefore, fertility treatments incorporated the use of second PGF2 $\alpha$  treatment 8 to 24 h after the PGF2 $\alpha$  of Ovsynch to enhance complete luteolysis rate and the chances of pregnancy following timed-AI.




Figure 2. Predicted probability of complete luteolysis based on concentrations of progesterone (P4 < 0.5 ng/ml 56, 72, and 96 h after PGF2 $\alpha$  injection) at time of PGF2 $\alpha$  injection of Ovsynch in lactating dairy cows with functional corpus luteum (CL; P4 concentrations  $\geq$ 0.24 ng/ml 24 h and  $\geq$ 0.09 ng/ml 56 h after treatment; n = 490) at time of treatment. Published in Journal of Dairy Science (Martins *et al.*, 2011a).

### Summary

It is critical to induce ovulation either during the estrous cycle or in an anovular condition to generate a new follicular wave and an accessory corpus luteum. Ovulation rate following a GnRH-induced LH surge is greatest on day 6 or 7 of the estrous cycle during the latter stages of the first follicular wave. It is also critical to initiate the induction of luteolysis of the spontaneous-formed CL as well as the accessory CL when the dominant follicle from the new follicular wave is at an ideal stage of maturity. Induction of complete luteolysis

of these corpora lutea is critical and cannot be jeopardized; therefore, it is imperative to utilize two doses of PGF2 $\alpha$  8 to 24 h apart. In essence, inducing the initiation of a new wave and causing ovulation of the DF from that new wave manipulates the age of the ovulatory follicle similar to that of heifers during an estrous cycle (Sartori *et al.*, 2002, 2004; Wolfenson *et al.*, 2004). Presynch-10 or 11, G6G, and Double Ovsynch create these physiological differences at a greater rate compared to Ovsynch alone and Presynch 12 or 14 and may increase pregnancies per AI outcomes 30 to 60% (Bello *et al.*, 2006; Galvão *et al.*, 2007;



Souza *et al.*, 2008; Ribeiro *et al.*, 2012; Astiz and Fargas, 2013; Dirandeh *et al.*, 2015).

### References

- Astiz S, Fargas O.** 2013. Pregnancy per AI differences between primiparous and multiparous high-yield dairy cows after using Double Ovsynch or G6G synchronization protocols. *Theriogenology*, 79:1065-1070.
- Bello NM, Steibel JP, Pursley JR.** 2006. Optimizing ovulation to first GnRH improved outcomes to each hormonal injection of ovsynch in lactating dairy cows. *J Dairy Sci*, 89:3413-3424
- Bello NM, Pursley JR.** 2007. Strategies to maximize ovulation to first GnRH of Ovsynch in lactating dairy cows. *J Dairy Sci*, 90:326-327. (abstract).
- Bergfeld EG, Kojima FN, Cupp AS, Wehrman ME, Peters KE, Mariscal V, Sanchez T, Kinder JE.** 1996. Changing dose of progesterone results in sudden changes in frequency of luteinizing hormone pulses and secretion of 17 beta-estradiol in bovine females. *Biol Reprod*, 54:546-553.
- Bisinotto RS, Ribeiro ES, Martins LT, Marsola RS, Greco LF, Favoreto MG, Risco CA, Thatcher WW, Santos JEP.** 2010. Effect of interval between induction of ovulation and artificial insemination (AI) and supplemental progesterone for resynchronization on fertility of dairy cows subjected to a 5-d timed AI program. *J Dairy Sci*, 93:5798-5808.
- Brusveen DJ, Cunha AP, Silva CD, Cunha PM, Sterry RA, Silva EPB, Guenther JN, Wiltbank MC.** 2008. Altering the time of the second gonadotropin-releasing hormone injection and artificial insemination (AI) during Ovsynch affects pregnancies per AI in lactating dairy cows. *J Dairy Sci*, 91:1044-1052
- Brusveen DJ, Souza AH, Wiltbank MC.** 2009. Effects of additional prostaglandin F2alpha and estradiol-17beta during Ovsynch in lactating dairy cows. *J Dairy Sci*, 92:1412-1422.
- Butler WR, Smith RD.** 1989. Interrelationships between energy balance and postpartum reproductive function in dairy cattle. *J Dairy Sci*, 72:767-783.
- Dirandeh E, Roodbari AR, Colazo MG.** 2015. Double-Ovsynch, compared with presynch with or without GnRH, improves fertility in heat-stressed lactating dairy cows. *Theriogenology*, 83:438-443.
- Escalante RC, Poock SE, Mathew DJ, Martin WR, Newsom EM, Hamilton SA, Pohler KG, Lucy MC.** 2013. Reproduction in grazing dairy cows treated with 14-day controlled internal drug release for presynchronization before timed artificial insemination compared with artificial insemination after observed estrus. *J Dairy Sci*, 96:300-306.
- Fricke PM, Giordano JO, Valenza A, Lopes G, Amundson MC, Carvalho PD.** 2014. Reproductive performance of lactating dairy cows managed for first service using timed artificial insemination with or without detection of estrus using an activity-monitoring system. *J Dairy Sci*, 97:2771-2781.
- Galvão KN, Sá Filho MF, Santos JEP.** 2007. Reducing the interval from presynchronization to initiation of timed artificial insemination improves fertility in dairy cows. *J Dairy Sci*, 90:4212-4218.
- Giordano JO, Wiltbank MC, Guenther JN, Pawlisch R, Bas S, Cunha AP, Fricke PM.** 2012. Increased fertility in lactating dairy cows resynchronized with Double-Ovsynch compared with Ovsynch initiated 32 d after timed artificial insemination. *J Dairy Sci*, 95:639-653.
- Giordano JO, Wiltbank MC, Fricke PM, Bas S, Pawlisch R, Guenther JN, Nascimento AB.** 2013. Effect of increasing GnRH and PGF2 $\alpha$  dose during Double-Ovsynch on ovulatory response, luteal regression, and fertility of lactating dairy cows. *Theriogenology*, 80:773-783.
- Her E, Wolfenson D, Flamenbaum I, Folman Y, Kaim M, Berman A.** 1988. Thermal, productive, and reproductive responses of high yielding cows exposed to short-term cooling in summer. *J Dairy Sci*, 71:1085-1092.
- Herlihy MM, Giordano JO, Souza AH, Ayres H, Ferreira RM, Keskin A, Nascimento AB, Guenther JN, Gaska JM, Kacuba SJ, Crowe MA, Butler ST, Wiltbank MC.** 2012. Presynchronization with Double-Ovsynch improves fertility at first postpartum artificial insemination in lactating dairy cows. *J Dairy Sci*, 95:7003-7014.
- Lopez H, Caraviello DZ, Satter LD, Fricke PM, Wiltbank MC.** 2005. Relationship between level of milk production and multiple ovulations in lactating dairy cows. *J Dairy Sci*, 88:2783-2793.
- López-Gatius F, Santolaria P, Yániz J, Rutlant J, López-Béjar M.** 2002. Factors affecting pregnancy loss from gestation day 38 to 90 in lactating dairy cows from a single herd. *Theriogenology*, 57:1251-1261.
- López-Gatius F, Hunter RHF.** 2005. Spontaneous reduction of advanced twin embryos: its occurrence and clinical relevance in dairy cattle. *Theriogenology*, 63:118-125.
- Lucy MC.** 2001. Reproductive loss in high-producing dairy cattle: where will it end? *J Dairy Sci*, 84:1277-1293.
- Martin AD, Kielland C, Nelson ST, Østerås O.** 2015. The effects of building design on hazard of first service in Norwegian dairy cows. *J Dairy Sci*, 98:8655-8663.
- Martins JPN, Policelli RK, Neuder LM, Raphael W, Pursley JR.** 2011a. Effects of cloprostenol sodium at final prostaglandin F2 $\alpha$  of Ovsynch on complete luteolysis and pregnancy per artificial insemination in lactating dairy cows. *J Dairy Sci*, 94:2815-2824.
- Martins JPN, Policelli RK, Pursley JR.** 2011b. Luteolytic effects of cloprostenol sodium in lactating dairy cows treated with G6G/Ovsynch. *J Dairy Sci*, 94:2806-2814.
- Orihuela A.** 2000. Some factors affecting the behavioural manifestation of oestrus in cattle: a review. *Appl Anim Behav Sci*, 70:1-16.
- Platz S, Ahrens F, Bendel J, Meyer HHD, Erhard MH.** 2008. What happens with cow behavior when replacing concrete slatted floor by rubber coating: a case study. *J Dairy Sci*, 91:999-1004.
- Pursley JR, Mee MO, Wiltbank MC.** 1995. Synchronization of ovulation in dairy cows using



- PGF2alpha and GnRH. *Theriogenology*, 44:915-923.
- Pursley JR, Kosorok MR, Wiltbank MC.** 1997a. Reproductive management of lactating dairy cows using synchronization of ovulation. *J Dairy Sci*, 80:301-306.
- Pursley JR, Wiltbank MC, Stevenson JS, Ottobre JS, Garverick HA, Anderson LL.** 1997b. Pregnancy rates per artificial insemination for cows and heifers inseminated at a synchronized ovulation or synchronized estrus. *J Dairy Sci*, 80:295-300.
- Pursley JR, Martins JPN.** 2011. Impact of circulating concentrations of progesterone and antral age of the ovulatory follicle on fertility of high-producing lactating dairy cows. *Reprod Fertil Dev*, 24:267-271.
- Ribeiro ES, Monteiro APA, Lima FS, Ayres H, Bisinotto RS, Favoreto M, Greco LF, Marsola RS, Thatcher WW, Santos JEP.** 2012. Effects of presynchronization and length of proestrus on fertility of grazing dairy cows subjected to a 5-day timed artificial insemination protocol. *J Dairy Sci*, 95:2513-2522.
- Sartori R, Fricke PM, Ferreira JC, Ginther OJ, Wiltbank MC.** 2001. Follicular deviation and acquisition of ovulatory capacity in bovine follicles. *Biol Reprod*, 65:1403-1409.
- Sartori R, Rosa GJM, Wiltbank MC.** 2002. Ovarian structures and circulating steroids in heifers and lactating cows in summer and lactating and dry cows in winter. *J Dairy Sci*, 85:2813-2822.
- Sartori R, Haughian JM, Shaver RD, Rosa GJM, Wiltbank MC.** 2004. Comparison of ovarian function and circulating steroids in estrous cycles of Holstein heifers and lactating cows. *J Dairy Sci*, 87:905-920.
- Savio JD, Boland MP, Roche JF.** 1990. Development of dominant follicles and length of ovarian cycles in post-partum dairy cows. *J Reprod Fertil*, 88:581-591.
- Souza AH, Gumen A, Silva EPB, Cunha AP, Guenther JN, Peto CM, Caraviello DZ, Wiltbank MC.** 2007. Supplementation with Estradiol-17 $\beta$  before the last gonadotropin-releasing hormone injection of the ovsynch protocol in lactating dairy cows. *J Dairy Sci*, 90:4623-4634.
- Souza AH, Ayres H, Ferreira RM, Wiltbank MC.** 2008. A new presynchronization system (Double-Ovsynch) increases fertility at first postpartum timed AI in lactating dairy cows. *Theriogenology*, 70:208-215.
- Spalding RW, Everett RW, Foote RH.** 1975. Fertility in New York artificially inseminated Holstein herds in dairy herd improvement. *J. Dairy Sci*, 58:718-723.
- Strickland J, Martins JPN, Neuder LM, Pursley JR.** 2010. Effect of 14/11 Presynch/Ovsynch on 1st service conception rates of lactating dairy cows compared to AI following a detected estrus. In: Abstracts of the 43rd Annual Conference of the American Association of Bovine Practitioners, 2010, Albuquerque, NM. Auburn, AL: AABP.
- Vasconcelos JLM, Silcox RW, Rosa GJ, Pursley JR, Wiltbank MC.** 1999. Synchronization rate, size of the ovulatory follicle, and pregnancy rate after synchronization of ovulation beginning on different days of the estrous cycle in lactating dairy cows. *Theriogenology*, 52:1067-1078.
- Vasconcelos JLM, Sangsritavong S, Tsai SJ, Wiltbank MC.** 2003. Acute reduction in serum progesterone concentrations after feed intake in dairy cows. *Theriogenology*, 60:795-807.
- Washburn SP, Silvia WJ, Brown CH, McDaniel BT, McAllister AJ.** 2002. Trends in reproductive performance in Southeastern Holstein and Jersey DHI herds. *J Dairy Sci*, 85:244-251.
- Wiltbank MC, Fricke PM, Sangsritavong S, Sartori R, Ginther OJ.** 2000. Mechanisms that prevent and produce double ovulations in dairy cattle. *J Dairy Sci*, 83:2998-3007.
- Wiltbank M, Lopez H, Sartori R, Sangsritavong S, Gümen A.** 2006. Changes in reproductive physiology of lactating dairy cows due to elevated steroid metabolism. *Theriogenology*, 65:17-29.
- Wiltbank MC, Souza AH, Carvalho PD, RW. Bender, Nascimento AB.** 2011. Improving fertility to timed artificial insemination by manipulation of circulating progesterone concentrations in lactating dairy cattle. *Reprod Fertil Dev*, 24:238-243.
- Wiltbank MC, Pursley JR.** 2014. The cow as an induced ovulator: timed AI after synchronization of ovulation. *Theriogenology*, 81:170-185.
- Wolfenson D, Inbar G, Roth Z, Kaim M, Bloch A, Braw-Tal R.** 2004. Follicular dynamics and concentrations of steroids and gonadotropins in lactating cows and nulliparous heifers. *Theriogenology*, 62:1042-1055.
- Xu Z, Garverick HA, Smith GW, Smith MF, Hamilton SA, Youngquist RS.** 1995. Expression of follicle-stimulating hormone and luteinizing hormone receptor messenger ribonucleic acids in bovine follicles during the first follicular wave. *Biol Reprod*, 53:951-957.
-